

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C12N 15/11, 15/70, 1/21, 1/12, 15/82, A01H 5/00	A2	(11) International Publication Number: WO 97/04089 (43) International Publication Date: 6 February 1997 (06.02.97)
(21) International Application Number: PCT/US96/11999 (22) International Filing Date: 19 July 1996 (19.07.96) (30) Priority Data: PCT/US95/09098 20 July 1995 (20.07.95) WO (34) Countries for which the regional or international application was filed: US et al. (71) Applicants (for all designated States except US): SUMITOMO CHEMICAL COMPANY, LTD. [JP/JP]; 5-33, Kitahama 4-chome, Chuo-ku, Osaka 541 (JP). DUKE UNIVERSITY [US/US]; 012 Allen Building, Durham, NC 27708 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): SATO, Ryo [JP/JP]; 2-10-5-301 Sonehigashinocho, Toyonaka-shi, Osaka 561 (JP). BOYNTON, John [US/US]; 1808 Woodburn Road, Durham, NC 27705 (US). GILLHAM, Nicholas, W. [US/US]; 1211 Woodburn Road, Durham, NC 27705 (US). HARRIS, Elizabeth, H. [US/US]; 3413 Rolling Hill Road, Durham, NC 27705 (US). (74) Agents: BIRCH, Anthony, L. et al.; Birch, Stewart, Kolasch & Birch, L.L.P., P.O. Box 747, Falls Church, VA 22040-0747 (US).	(81) Designated States: CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(54) Title: PORPHYRIN-ACCUMULATING TYPE HERBICIDE RESISTANCE GENE (57) Abstract Provided are DNA fragments and biologically functional equivalents thereof that confer resistance to porphyrin-accumulating type herbicides upon plant and algal cells, plasmids containing these DNA fragments or biologically functional equivalents thereof, microorganisms containing these DNA fragments or biologically functional equivalents thereof, methods for conferring resistance to porphyrin-accumulating type herbicides upon plant or algal cells using these DNA fragments or biologically functional equivalents thereof, and herbicide-resistant plants or algae into which these DNA fragments or biologically functional equivalents thereof have been introduced and in which they are expressed.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LÜ	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

Porphyrin-Accumulating Type Herbicide Resistance Gene

Background of the Invention

Related Applications

The present application is a Continuation-In-Part
5 of international application PCT/US95/09098, filed July
20, 1995.

Field of the Invention

The present invention relates to DNA fragments that
confer resistance to porphyrin-accumulating type
10 herbicides on plant and algal cells, plasmids and
microorganisms that contain these DNA fragments, methods
for conferring resistance to porphyrin-accumulating type
herbicides onto plant and algal cells by using these DNA
fragments, and plants and algae into which these DNA
15 fragments have been introduced for the purpose of
conferring resistance to such herbicides thereon.

Description of Related Art

A group of widely-known compounds used as active
ingredients of some varieties of commercially- and
20 otherwise-available herbicides exhibit herbicidal
activity in the presence of light, but exhibit no
herbicidal activity in darkness. This has led to their
common designation as light-dependent or porphyrinic
herbicides. It has recently been shown that these
25 herbicides induce high levels of porphyrin accumulation

in plants and algae, and thus they are now designated as "porphyrin-accumulating type herbicides" [Zoku, Iyaku hin-no-Kaihatsu, (translation: "The Development of Medical Drug Products; continuation") vol. 18; Development of Agricultural Chemicals II, chapter 16, section 16 - 1, Hajime Iwamura, Tamio Ueno & Katsuzo Kamoshita, eds., Hirokawa Shoten, Tokyo, pubs.) or simply "porphyric herbicides". It was reported by Matringe, M., Camadro, J.M., Labbe, P. & Scalla, R. (Biochem J. 260:231 (1989)) and by Matringe, M., Camadro, J.M., Labbe, P. & Scalla, R. (FEBS Lett. 245:35 (1989)) that porphyrin-accumulating type herbicides (referred to below also as porphyric herbicides) inhibit isolated protoporphyrinogen oxidase (referred to below as "protox").

Since most crop plants do not exhibit resistance to these porphyric herbicides, it is not possible to use these herbicides on farmland when such crops are under cultivation. If it were possible to develop crops resistant to porphyric herbicides, such herbicides could be used on these crops. This would make crop management easier, and increase the value of these herbicides in agricultural applications. For this reason, it is desirable to develop a method for conferring resistance to porphyrin-accumulating type herbicides upon crop plants.

Summary of the Invention

With this goal in mind, the present inventors have investigated a mutant strain, designated RS-3, of the unicellular green alga *Chlamydomonas reinhardtii* which displays specific resistance to porphyric herbicides. Wild-type strains of this alga are normally highly sensitive to porphyric herbicides. The present inventors have discovered that inhibition by porphyric herbicides of protox activity in chloroplast fragments isolated from the RS-3 strain of *Chlamydomonas reinhardtii* was significantly lower than in chloroplast

fragments from the wild-type strain. The inventors therefore constructed a genomic DNA library from total nuclear DNA isolated from the RS-3 mutant strain, and succeeded in isolating clones that contain a gene
5 responsible for resistance to porphyrin herbicides. Thus, the inventors were able to obtain DNA fragments that can confer resistance to porphyrin-accumulating type herbicides onto plant and algal cells.

Accordingly, it is an object of the present
10 invention to provide an isolated, purified DNA fragment that confers resistance to porphyrin-accumulating type herbicides when expressed in plant or algal cells, plasmids and microorganisms containing said DNA fragment. A DNA fragment according to the present
15 invention preferably has a nucleotide sequence of one or more portions of DNA comprising the genome of an alga, or has a nucleotide sequence highly homologous to the nucleotide sequence of DNA comprising one or more portions of the genome of an alga.

20 Additional objects of the present invention are a method for conferring resistance to porphyrin-accumulating type herbicides upon plant or algal cells, comprising introducing said DNA fragment into said plant or algal cells, wherein said DNA fragment is expressed;
25 and plants or algae into which said DNA fragment has been introduced, wherein said DNA fragment is expressed, thereby conferring herbicide resistance upon said plants or algae.

Another object of the present invention is to
30 provide an isolated, purified DNA fragment having the following characteristics:

a) comprising a nucleotide sequence derived from a DNA fragment obtained from a strain of the unicellular green alga *Chlamydomonas reinhardtii* that exhibits
35 resistance to porphyrin-accumulating type herbicides;

b) containing restriction sites for XhoI, PstI, PstI, PstI, PstI, BamHI, SalI, SalI, and XhoI, and

having a restriction site map as shown in Figure 1(a);

c) having a molecular size of approximately 3.4 kb;

and

d) which confers resistance to porphyrin-

5 accumulating type herbicides in plant or algal cells when expressed therein, or a biologically functional equivalent thereof.

A further object of the present invention is to provide an isolated, purified DNA fragment having the
10 following characteristics:

a) comprising a nucleotide sequence derived from a DNA fragment obtained from a strain of the unicellular green alga *Chlamydomonas reinhardtii* that exhibits resistance to porphyrin-accumulating type herbicides;

15 b) containing restriction sites for EcoRI, XhoI, PstI, PstI, PstI, PstI, PstI, BamHI, SalI, SalI, XhoI and HindIII, and having a restriction site map as shown in Figure 1(b);

c) having a molecular size of approximately 9.9 kb;

20 and

d) which confers resistance to porphyrin-accumulating type herbicides in plant or algal cells when expressed therein, or a biologically functional equivalent thereof.

25 Another object of the present invention is to provide an isolated, purified DNA fragment having the following characteristics:

a) comprising a nucleotide sequence derived from a DNA fragment obtained from a strain of the unicellular
30 green alga *Chlamydomonas reinhardtii* that exhibits resistance to porphyrin-accumulating type herbicides;

b) containing restriction sites for EcoRI, XhoI, PstI, PstI, PstI, PstI, PstI, BamHI, SalI, SalI, XhoI, HindIII, and KpnI, and having a restriction site map as
35 shown in Figure 1(c);

c) having a molecular size of approximately 10.0 kb; and

d) which confers resistance to porphyrin-accumulating type herbicides in plant or algal cells when expressed therein, or a biologically functional equivalent thereof.

5 A further object of the present invention is to provide an isolated, purified DNA fragment having the following characteristics:

a) comprising a nucleotide sequence derived from a DNA fragment obtained from a strain of the unicellular
10 green alga *Chlamydomonas reinhardtii* that exhibits resistance to porphyrin-accumulating type herbicides;

b) containing restriction sites for EcoRI, XhoI, PstI, PstI, PstI, PstI, PstI, BamHI, SalI, SalI, XhoI, HindIII, BamHI, SalI, HindIII, and KpnI, and having a
15 restriction site map as shown in Figure 1(d);

c) having a molecular size of approximately 13.8 kb; and

d) which confers resistance to porphyrin-accumulating type herbicides in plant or algal cells
20 when expressed therein, or a biologically functional equivalent thereof.

Further objects of the present invention are to provide plasmids and microorganisms containing any of the foregoing DNA fragments or biologically functional
25 equivalents thereof, a method of conferring resistance to porphyrin-accumulating type herbicides upon plant or algal cells, comprising introducing said DNA fragments or biologically functional equivalents thereof into plant or algal cells in a functionally operable manner
30 so that said DNA fragments or biologically functional equivalents thereof are expressed in said plant or algal cells, and the expression of the DNA fragment confers resistance to porphyrin-accumulating type herbicides upon the transformed plant or algal cells. It is
35 preferred that cells cultured in vitro that have been transformed by the DNA fragments of the invention in a functionally operable manner are resistant to a

porphyrin-accumulating type herbicide at a concentration of at least 0.01 μM , preferably at a concentration of at least 0.03 μM , most preferably at a concentration of at least 0.1 μM herbicide. When compound A or compound B are used as the test compounds, the range of concentration is preferably 0.01 to 0.3 μM , more preferably 0.03 to 0.6 μM , most preferably 0.1 to 0.3 μM . Otherwise the range is between 0.01 to 30 μM , more preferably 0.03 to 10 μM , most preferably 0.1 to 3 μM . The concentration of herbicide used to test resistance of transformed plants or tissues therefrom is to the high end of these ranges or even higher, and can be determined by the ordinarily skilled artisan by experimentation typical in the art. The herbicide used for testing herbicide resistance of cells in vitro or of whole transformed plants or algae is preferably a N-phenyl-tetrahydrophthalimide compound. N-(4-chloro-2-fluoro-5-propargyloxy)phenyl-3,4,5,6-tetrahydrophthalimide (compound A) or 7-fluoro-6-[(3,4,5,6)-tetrahydrophthalimido]-4-(2-propynyl)-1,4-benzoxazin-3(2H)-one (referred to below as "compound B") are especially preferred for this purpose.

Another object of the present invention is to provide plants or algae into which have been introduced in a functionally operable manner said DNA fragments or biologically functional equivalents thereof.

A still further object of the present invention is to provide an isolated, purified genomic DNA fragment comprising the nucleotide sequence shown in SEQ ID NO:1; plasmids and microorganisms containing said DNA fragment; a method of conferring resistance to porphyrin-accumulating type herbicides upon plant or algal cells, comprising introducing the cDNA corresponding to the mRNA encoded by said DNA fragment that confers porphyrin herbicide resistance on plant or algal cells, wherein said cDNA is expressed; and plants

or algae into which cDNA corresponding to the mRNA encoded by said DNA fragment having the nucleotide sequence shown in SEQ ID NO:1 has been introduced, wherein said cDNA is expressed.

5 Yet further objects of the present invention include the use of any of the DNA fragments or biologically functional equivalents thereof disclosed herein as a genetic marker (for herbicide resistance), to produce a recombinant plasmid or transformed
10 microorganism, to produce a probe useful in identifying related DNA sequences that confer resistance to porphyrin-accumulating type herbicides in plant and algal cells, and to produce plants or algae resistant to porphyrin-accumulating type herbicides.

15 The DNA fragments and biologically functional equivalents thereof of the present invention are hereinafter referred to as the "subject nucleic acid fragments" or "subject DNA fragments". Specific individual fragments will be designated by their
20 restriction sites and molecular sizes (kb).

The present invention includes plasmids containing the above-mentioned DNA fragments or their biologically functional equivalents (hereinafter referred to as the "subject plasmids"), microorganisms containing these DNA
25 fragments or their equivalents (hereinafter referred to as the "subject microorganisms"), plants or algae containing these DNA fragments or their equivalents (hereinafter referred to as the "subject plants"), and methods for conferring resistance to porphyrin-accumulating type herbicides upon plant and algal cells
30 by using these DNA fragments or their equivalents.

With regard to the terminology used herein, the term "DNA fragments" refers not only to the subject DNA fragments, but also to degenerate isomers and genetical-
35 ly equivalent modified forms of these fragments. "Degenerate isomers" is taken here to mean isomers whose nucleotide base sequence is degenerately related to the

original fragments; that is, all nucleic acid fragments including the corresponding mRNA or corresponding cDNA, that contain essentially the same genetic information as the original fragments. "Genetically equivalent modified forms" is taken here to mean DNA fragments that may have undergone base changes, additions, or deletions, but which essentially contain the same inherent genetic information as the original fragments.

Specific examples of the latter include DNA fragments whose nucleotide sequence shows high homology to the subject nucleic acid fragments, which are readily isolated using conventional DNA-DNA or DNA-RNA hybridization techniques, or that are amplified using known PCR (Polymerase Chain Reaction) methods, and which possess the ability to confer resistance to porphyrin-accumulating type herbicides when introduced by conventional transformation techniques into plant or algal cells normally sensitive to these herbicides.

The phrase "porphyrin-accumulating type herbicide" or the phrase "porphyrin herbicides" refers to light-dependent herbicides, i.e., compounds that kill sensitive plants in the presence of light, but which exhibit no herbicidal activity in darkness, and which induce the accumulation of high levels of porphyrins in plants to which they have been applied. These herbicides include, for example, oxadiazon, flupropacil, [N-(4-chloro-2-fluoro-5-propargyloxy)phenyl-3,4,5,6-tetrahydrophtalimide (referred to below as compound A), the diphenyl ether herbicides such as acifluorfen, lactofen, oxyfluorfen, as well as the following: pentyl [2-chloro-5-(cyclohex-1-ene-1,2-dicarboximido)-4-fluor-phenoxy] acetate, 7-fluoro-6-[(3,4,5,6)-tetrahydrophthalimido]-4-(2-propynyl)-1,4-benzoxazin-3(2H)-one (referred to below as "compound B"), 6-[(3,4,5,6-tetrahydro)phthalimido]-4-(2-propynyl)-1,4-benzoxazin-3(2H)-one, 2-[7-fluoro-3-oxo-4-(2-propynyl)-3,4-dihydro-2H-1,4-benzox-

azin-6-yl]perhydroimidazo[1,5-a]pyridine-1,3-dione,
2-[(4-chloro-2-fluoro-5-propargyloxy)phenyl] perhydro-1-
H-1,2,4-triazolo-[1,2-a]pyridazine-1,3-dione, 2-[7-fluo-
ro-3-oxo-4-(2-propynyl)-3,4-dihydro-2H-1,4-benzoxazin-
5 6-yl]5,6,7,8-1,2,4-triazolo[4,3-a]pyridine-3H-one,
2-[3-oxo-4-(2-propynyl)-3,4-dihydro-2H-1,4-benzoxazin-
6-yl]-1-methyl-6-trifluoromethyl-2,4(1H,3H)-pyrimidine
dione, 2-[6-fluoro-2-oxo-3-(2-propynyl)-2,3-
dihydrobenzthiazol-5-yl]-3,4,5,6-tetrahydrophthalimide,
10 1-amino-2-[3-oxo-4-(2-propynyl)-3,4-dihydro-2H-1,4-
benzoxazin-6-yl]-6-tri-fluoromethyl-2,4(1H,3H)-
pyrimidinedione, as well as analogs of these compounds.

The subject nucleic acid fragments may be con-
structed by the artificial synthesis of their nucleotide
15 sequences; however, they are more typically isolated
from a mutant strain of the unicellular green alga
Chlamydomonas reinhardtii, designated RS-3, that is
resistant to porphyrin-accumulating type herbicides.
Said mutant strain RS-3 is stored at the Chlamydomonas
20 Genetics Center (address: DCMB Group, Department of
Botany, Box 91000, Duke University, Durham, NC,
27708-1000, USA) under the entry number CC-2674. Thus,
the mutant strain RS-3 is publicly available for
distribution. As will be described below, the
25 microorganisms that host the plasmids containing the
subject nucleic acid fragments are also on deposit under
the terms of the Budapest Treaty, and are thus freely
available as well. The plasmids hosted by these
microorganisms can be readily extracted using
30 conventional techniques and the subject fragments
recovered by reference to the restriction maps shown in
Figures 1(a)-1(d). It would be possible, for example,
to introduce specific alterations into these fragments
using PCR or other site-directed mutagenesis techniques,
35 or to use the subject nucleic acid fragments or their
corresponding cDNAs, PCR products, or oligonucleotides
as probes to isolate other DNA fragments exhibiting high

homology to the subject nucleic acid fragments, and thus to generate homologs as discussed above.

Further scope of the applicability of the present invention will become apparent from the detailed description and drawings provided below. It should be understood, however, that the following detailed description and specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

Brief Description of the Drawings

The above and other objects, features, and advantages of the present invention will be better understood from the following detailed descriptions taken in conjunction with the accompanying drawings, all of which are given by way of illustration only and are not limitative of the present invention, in which:

Figure 1(a)-1(d) shows restriction site maps of cloned DNA fragments of various sizes that confer resistance to porphyrin-accumulating type herbicides. The sizes of the fragments are indicated by the numbers (kb) in Figure 1(e). Abbreviations: B, BamHI; S, SalI; P, PstI; X, XhoI; E, EcoRI; H, HindIII; K, KpnI; C, ClaI.

Figure 1(a): 3.4 kb DNA fragment designated as Xho3.4;

Figure 1(b): 9.9 kb DNA fragment designated as Hind9.9;

Figure 1(c): 10.0 kb DNA fragment designated as Hind10.0;

Figure 1(d): 13.8 kb DNA fragment designated as Eco13.8;

Figure 1(e): approximately 40 kb DNA fragment harbored by cosmid clone 2955 (Cos2955).

Figure 2 shows the structure of a pBS plasmid containing an Xho3.4 fragment insert. Distances between restriction sites (kb) are shown by the numbers in the insert.

5 Figure 3 shows the structure of a pBS plasmid containing a Hind10.0 fragment insert. Distances between restriction sites (kb) are shown by the numbers in the insert. Figures 1(c) and 2 show the PstI sites.

10 Figure 4 shows the structure of a pBS plasmid containing an Eco13.8 fragment insert. Distances between restriction sites (kb) are shown by the numbers in the insert. Figures 1(d) and 2 show the PstI sites.

Detailed Description of the Invention

15 The following detailed description of the invention is provided to aid those skilled in the art in practicing the present invention. Even so, the following detailed description should not be construed to unduly limit the present invention, as modifications and variations in the embodiments herein discussed may
20 be made by those of ordinary skill in the art without departing from the spirit or scope of the present inventive discovery.

 The contents of each of the references cited herein are herein incorporated by reference in their entirety.

25 Overview

 The present nucleic acid fragments are obtained by conventional genetic engineering protocols as described in publications such as Molecular Cloning, 2nd Edition, by J. Sambrook, E. F. Frisch, and T. Maniatis, Cold
30 Spring Harbor Publications (1989). Specifically, genomic DNA is extracted from the mutant strain RS-3 according to a protocol such as that described by E.H. Harris, The Chlamydomonas Source Book, pp. 610-613 (Chapter 12), Academic Press, San Diego (1989). Namely,
35 *C. reinhardtii* cells are lysed and the DNA is extracted

by treatment with a protease and surface active agents such as SDS or Sarkosyl. Genomic DNA is subsequently extracted by conventional techniques involving phenol-chloroform extraction, centrifugation, etc., to
5 remove proteins, after which the DNA is recovered by ethanol precipitation. The DNA thus obtained can be further purified by sodium iodide-ethidium bromide density gradient centrifugation, and the lowermost, major band corresponding to nuclear genomic DNA
10 recovered. Nuclear genomic DNA thus obtained is partially digested using an appropriate restriction enzyme such as Sau3AI. Linkers or adaptors are attached to both ends of the DNA fragments thus obtained using T4 DNA ligase. If necessary, excess free linkers or adapt-
15 ors can be removed by gel filtration, and the fragments can then be inserted into an appropriate commercially available cosmid vector or a phage vector such as those derived from λ phage. Phage particles generated by *in vitro* packaging are transfected into *E. coli* and allowed
20 to form colonies or plaques on solid media. A genomic DNA library can be obtained by isolating and maintaining individual *E. coli* clones harboring hybrid cosmids or by conventional methods for isolating and maintaining *E. coli* clones or phage particles in a mixture.

25 Since no porphyric herbicide resistance gene had been previously isolated and characterized from any plant or algal species prior to the present invention, it was not feasible to screen the genomic DNA library described above by synthesis of an oligonucleotide probe
30 corresponding to the deduced nucleotide sequence of such a gene, labeling this probe with a radioisotope or fluorescent tag, and using this to select genomic DNA clones containing the subject DNA fragments. Therefore, the genomic clones containing subject DNA fragments were
35 screened by transforming a strain of *Chlamydomonas reinhardtii* sensitive to porphyric herbicides with the genomic DNA from the cosmid library using normal trans-

formation techniques for this organism (Kindle, K., Proc. Natl. Acad. Sci. USA Vol. 87, p. 1228 (1990); Boynton J.E. and Gillham, N.W., Methods In Enzymology: Recombinant DNA, Part H, R. Wu, Ed., Academic Press, San Diego, CA, Vol. 217, p. 510, (1993)) to isolate hybrid cosmids containing nuclear genomic DNA fragments capable of conferring resistance to a porphyric herbicide. A restriction map of the hybrid cosmid clone thus obtained was determined, various restriction fragments were subcloned into the pBluescript vector, and subclones that conferred resistance to porphyric herbicides onto normally sensitive *Chlamydomonas* strains were selected. Using the subject DNA fragments and the subject plasmids as starting material, the nucleotide sequence of the 3.4 kb fragment was determined by the method of Sanger (Sanger, F. and Coulson, A.R. J. Mol. Biol., Vol. 94, p. 441 (1975); Sanger, F., Nicklen, and Coulson, A.R. Proc. Natl. Acad. Sci. USA, Vol. 74, p. 5463 (1977)) or improved versions of this method. Sequences of the larger 9.9, 10.0, and 13.8 kb fragments or the cDNAs corresponding to these fragments can be determined by the same methodology. The transcriptional initiation site of the porphyric herbicide resistance gene can be localized in one or more of these overlapping fragments using the primer extension technique described by Bina-Stem, M. et al. (Proc. Natl. Acad. Sci. USA, Vol. 76, p. 731, (1977)) and Sollner-Webb and Reeder, R.H. (Cell, Vol. 18, p. 485 (1978)), or by the S1 mapping technique described by Berk, A.J. and Sharp, P.A. (Proc. Natl. Acad. Sci. USA, Vol. 75, p. 1274 (1978)). Typically, promoter sequences that are responsible for the regulation of gene expression are found in a region approximately 1 kb to 10 kb upstream of the transcription initiation site. The promoter region of the gene which confers resistance to porphyric herbicides can be determined by using standard *Chlamydomonas* transformation techniques (Kindle, K.,

Proc. Natl. Acad. Sci. USA, Vol. 87, p. 1228 (1990)) and chimeric reporter constructs. For example, various lengths of the region upstream of the transcription start site can be joined to an appropriate heterologous reporter gene such as GUS or one encoding enzymatically-determined antibiotic resistance. By introducing these constructs into *Chlamydomonas reinhardtii* using transformation and monitoring reporter gene expression, the promoter region of the gene conferring resistance to porphyric herbicides can ultimately be determined. In addition, a transcription terminator sequence is expected to be present within one or more of the overlapping cloned genomic DNA fragments downstream of the poly-A addition signal found in the 3' non-coding region downstream of the stop codon.

Herbicide-resistant transformants were obtained from the 13.8 and 10.0 kb fragments from RS-3 at about 80-fold higher frequency than from the 3.4 kb fragment. This is consistent with the 13.8 and 10.0 kb fragments containing the entire coding sequence plus upstream and downstream regulatory elements and integrating non-homologously and randomly in the nuclear genome of the herbicide-sensitive recipient strain. In contrast, the low transformation frequency observed with the 3.4 kb fragment is consistent with this fragment containing only a portion of the RS-3 gene which must integrate into the herbicide-sensitive RS-3 gene of the recipient by homologous recombination to be expressed. Nuclear transformants of *Chlamydomonas reinhardtii* arise much more frequently by random non-homologous recombination than by homologous recombination as has been demonstrated by experiments with the nuclear *nit-1* gene by Sodeinde, O.A. and Kindle, K.L., (Proc. Natl. Acad. Sci. USA Vol. 90, p.9199 (1993)).

The foregoing will be described in detail in the Examples presented below, although the present invention is not limited to these Examples.

Example 1Construction of *Chlamydomonas reinhardtii*Genomic DNA Library

The porphyric herbicide-resistant mutant strain
5 RS-3 of the unicellular alga *Chlamydomonas reinhardtii*
(*Chlamydomonas* Genetics Center, strain GB-2674) was
cultured mixotrophically under 200 $\mu\text{E}/\text{m}^2/\text{sec}$
photosynthetically active radiation with shaking for 5
days in TAP liquid medium composed of 7 mM NH_4Cl , 0.4 mM
10 $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.34 mM $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 25 mM potassium phosphate,
0.5 mM Tris (pH 7.0), 1 ml/L Hutner trace elements, and
1 ml/L glacial acetic acid (Harris, E. H., The
Chlamydomonas Sourcebook, Academic Press, San Diego,
1989, pp. 576-77) also containing 0.03 μM compound A.
15 Six liters of culture containing cells in early station-
ary growth phase (7.6×10^6 cells/ml) were harvested.
Cells were collected by centrifugation (8,000xg, 10 min,
4°C), resuspended in 50 ml of TEN buffer composed of 10
mM Tris-HCl, 10 mM EDTA, 150 mM NaCl, pH 8.0,
20 recentrifuged, and resuspended again in 50 ml of TEN
buffer. To this were gently added 5 ml of 20% (w/v)
SDS, 5 ml of 20% Sarkosyl, and 4 ml of protease solution
composed of 5 g of protease (Boehringer Mannheim No.
165921), 10 ml of 1M Tris-HCl (pH 7.5), and 0.11 g of
25 CaCl_2 in a total volume of 100 ml of deionized distilled
water. This was mixed by slowly rotating the solution
in a bottle for 24 hr at 4°C. Sixty ml of phenol-CIA
(phenol pre-saturated with TEN buffer and mixed well
with an equal volume of a chloroform:isoamyl-alcohol,
30 24:1, v/v) were subsequently added, and the contents
were rotated in the same bottle at room temperature for
1 hr.

The contents were then separated by centrifugation
(15,000xg, 20 min, room temperature), the aqueous
35 (upper) phase was recovered and mixed with 2 volumes of
95% (v/v) ethanol gently but thoroughly, and the DNA
precipitated by placing the contents at -20°C overnight.

The resulting precipitate was recovered by centrifugation (1,500xg, 20 min, 4°C) and washed once with ice-cold 70% (v/v) ethanol. Excess ethanol was removed and the DNA precipitate was dried under nitrogen flow
5 for 5 min at room temperature.

The dried precipitate was subsequently dissolved in 60 ml of 10mM Tris (pH 7.5), and under dim light, the following were added: 8 ml of 10-fold concentrated TEN buffer, 0.4 ml of ethidium bromide solution (10 mg/ml),
10 9.8 ml of 10 mM Tris-HCl (pH 7.5), and 120 ml of sodium iodide (NaI)-saturated TEN buffer. The contents were mixed by gently inverting the container and 25 ml were dispensed into each of 8 centrifuge tubes. These were subjected to ultracentrifugation in a Beckman 70 Ti rotor (44,000 rpm, 40 hr, 20°C). After ultra-
15 centrifugation, the lowermost, major band consisting of nuclear DNA was visualized by long-wave UV illumination, and recovered by use of a large-gauge syringe. The DNA in this band was again subjected to ultracentrifugation in a Beckman 70 Ti rotor (44,000 rpm, 44 hr, 20°C). The
20 purified nuclear DNA band was recovered as above.

Ethidium bromide was extracted from the solution containing the recovered nuclear DNA by adding isoamyl alcohol saturated with 1-2 volumes of TEN buffer and
25 subsequently discarding the alcohol (upper) phase. After repeating this step three times, the nuclear DNA from which ethidium bromide had been removed was precipitated by the addition of 2.5 volumes of ice-cold ethanol. The precipitate recovered was washed twice in
30 ice-cold 95% (v/v) ethanol, redissolved in a small volume of 10mM Tris-HCl (pH7.5), and stored at -20°C. An aliquot of this sample was diluted 100-fold and the concentration and purity of the DNA were quantified by measuring the absorbance at 260 nm and 280 nm.

35 Twenty five μ g of the genomic DNA thus obtained were partially digested by reaction with 0.83 units of the restriction enzyme Sau3AI at 37°C for 15 min in 277

5 μ l of 10 mM Tris-HCl buffer (pH 7.5) containing 50 mM NaCl, 10 mM MgCl₂, and 1 mM dithiothreitol. The reaction mixture was extracted with an equal volume of phenol equilibrated with Tris buffer (pH 7.5) followed by an equal volume of chloroform. Three M ammonium acetate was added to give a final concentration of 0.4M, followed by the addition of 2 volumes of ice-cold ethanol. This solution was mixed thoroughly and a DNA precipitate formed following storage of the sample overnight at -20°C. The precipitate was recovered by centrifugation in a tabletop centrifuge (10,000 rpm, 10 min), washed in 70% (v/v) ethanol and re-centrifuged. The precipitate was then resuspended in 20 μ l TE buffer (composed of 10 mM Tris-HCl, 0.1 mM EDTA·2Na), and the DNA was dephosphorylated by the addition of 70 μ l of deionized distilled water, 10 μ l of 10-fold concentrated CIAP buffer composed of 0.5M Tris-HCl (pH 8.5) and 1 mM EDTA, and 1 unit of CIAP (Calf Intestinal Alkaline Phosphatase). The total volume of 100 μ l was incubated for 60 min at 37°C and the reaction halted by the addition of 3 μ l of 0.5M EDTA (pH 8.0) and heat-treatment for 10 min at 68°C. The DNA was subjected to phenol and chloroform extractions and precipitated by the addition of ethanol containing ammonium acetate as described above.

The precipitate was washed with 70% (v/v) ethanol and the recovered DNA redissolved in TE buffer to a final concentration of 0.5 μ g/ml. Subsequently, the commercially available cosmid vector SuperCos-1 (Stratagene Inc.) was prepared following the protocol outlined in the SuperCos-1 instruction manual provided by the manufacturer. The vector was digested with the restriction enzyme XbaI, dephosphorylated with CIAP, re-digested with the restriction enzyme BamHI, recovered by ethanol precipitation, and redissolved in TE buffer to a final concentration of 1 μ g/ml. Two point five μ g of the prepared genomic DNA fragments were ligated to 1

μ g of the prepared SuperCos-1 vector in 20 μ l of reaction buffer (composed of 1 mM ATP, 50 mM Tris-HCl (pH7.5), 7 mM $MgCl_2$, and 1 mM dithiothreitol) by the addition of 2 units of T4 DNA ligase and incubation at 4°C overnight. Point five μ g of the hybrid cosmids thus generated were then packaged into λ phage particles capable of infecting *E. coli* by the use of a commercial *in vitro* phage packaging kit (Gigapack II XL, Stratagene Inc.) following the protocol outlined in the instruction manual provided. λ phage particles harboring these hybrid cosmids were then transfected into the *E. coli* strain NM554 (Stratagene, Inc.) by the procedure described below, and these *E. coli* cells were allowed to form colonies on LB medium plates (10 g/L NaCl, 10 g/L Bacto-tryptone, 5 g/L yeast extract, pH 7.5, 1.5% (w/v) agar) containing 50 μ g/ml ampicillin. The transfection protocol is as follows: (1) a single colony of *E. coli* strain NM554 was inoculated into 50 ml of medium (5g/L NaCl, 10g/L Bacto-tryptone, pH 7.4, 0.2% (w/v) maltose, 10mM $MgSO_4$) and cultured by shaking vigorously overnight at 37°C; (2) cells were collected by centrifugation (4,000 rpm, 10 min, 4°C) and resuspended in 10 mM $MgSO_4$ to an OD_{600} of 0.5; (3) 25 μ l of this bacterial suspension were mixed with 25 μ l of a 1/20th dilution of the phage particle solution harboring hybrid cosmids prepared as described above. The phage were infected into *E. coli* by allowing the mixture to stand at room temperature for 30 min. Two hundred μ l of LB medium (10 g/L NaCl, 10 g/L Tryptone, 5 g/L yeast extract) were subsequently added and the suspension was incubated at 37°C for 1 hr to allow for the expression of ampicillin resistance. The suspension was then plated onto LB medium plates containing 50 μ g/ml ampicillin and colonies formed following incubation at 37°C overnight. The transformation efficiency of the ampicillin marker was $1.7 \pm 0.1 \times 10^5$ transformants/ μ g DNA. The *E. coli* colonies containing hybrid cosmids thus obtained were

individually picked with sterile toothpicks and transferred into microtiter plate wells (Falcon, 24-well) each containing 0.5 ml of LB medium containing 50 µg/ml ampicillin and incubated without shaking at 37°C for 24
5 hr. Ten thousand and eighty individual clones were thereby isolated in 420 microtiter plates. Then, 187.5 µl of medium were removed from each well and combined in pools of 8 clones each (1.5 ml total) into 1,260
10 microtubes. The bacteria in each microtube were pelleted by centrifugation (10,000 rpm, 5 min, room temperature) and subjected to DNA extraction. The bacteria remaining in the microtiter plates were frozen at -70°C following the addition of an equal volume of 30% (w/v) glycerol. These plates were subsequently
15 stored at -20°C.

Example 2

Screening of the Genomic DNA Library

The various experimental methods used to screen the genomic DNA library are described below (methods A, B,
20 and C).

A. DNA extraction

Extraction of cosmid DNA from *E. coli* harboring the genomic DNA library generated as described in Example 1, as well as extraction of the plasmid pARG7.8 (Debuchy,
25 R., Purton, S., Rochaix, R.D., EMBO J., vol. 8, p. 2803, (1989)), utilized as a transformation control, was performed by standard extraction methods (for example, J. Sambrook, E. F. Frisch, T. Maniatis, Molecular Cloning, 2nd edition, Cold Spring Harbor Publications, (1989),
30 Vol. I, pp. 1.38 - 1.39). A description of the specific protocol follows.

The bacterial pellet in each microtube was thoroughly suspended in 100 µl of Solution I (composed of 50 mM glucose, 25 mM Tris-HCl (pH8.0), 10 mM EDTA),
35 to which 200 µl of Solution II (composed of 0.2N NaOH,

1% (w/v) SDS) were added. Each microtube was capped, the contents were gently mixed by inverting the tube 5-6 times, and the tube was cooled by placing on ice. One hundred and fifty μ l of ice-cold Solution III (composed of 60 ml of 5M potassium acetate (pH 4.8), 11.5 ml of glacial acetic acid, and 28.5 ml of deionized, distilled water) were subsequently added, the contents were mixed well, and the tubes cooled on ice for 5 min. The tubes were then centrifuged in a tabletop centrifuge (10,000 rpm, 2 min, 4°C) and the supernatant recovered. An equal volume of phenol:chloroform (1:1, v/v, pH 7.5) was added to the recovered supernatant, the contents were thoroughly mixed by vortexing, and the tubes were again centrifuged in a tabletop centrifuge (10,000 rpm, 2 min, 4°C), and the supernatant recovered. After re-extracting with chloroform, 900 μ l of ethanol were added to the supernatant and mixed. The DNA was precipitated by cooling the tubes on ice and the precipitates were recovered by centrifugation in a tabletop centrifuge (12,000 x g, 2 min, 4°C). The precipitates were washed in 70% (w/v) ethanol and recovered again by centrifugation (12,000 x g, 2 min, 4°C). Excess ethanol was removed by opening the microtube caps and allowing the ethanol to evaporate at room temperature for 10 min. The precipitates thus recovered were redissolved in 50 μ l of TE buffer (composed of 10 mM Tris-HCl (pH 7.5), 0.1 mM EDTA·2Na) to solubilize the DNA.

B. Transformation by the glass bead method

The glass bead transformation protocol, when employed, followed that described by Kindle, K. (Proc. Natl. Acad. Sci USA, vol. 87, p. 1228, (1990)). The actual protocol employed is presented below.

First, the unicellular green alga *Chlamydomonas reinhardtii* strain CC-425 (arginine auxotroph arg-2, cell wall deficient cw-15) was cultured mixotrophically

for 2 days to a cell density of $1-2 \times 10^6$ cells/ml in TAP liquid medium composed of 7 mM NH_4Cl , 0.4 mM $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.34 mM $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 25 mM potassium phosphate-0.5 mM Tris (pH 7.0), 1 ml/L Hutner trace elements, 1 ml/L glacial acetic acid (described in Harris, E. H., The Chlamydomonas Sourcebook, Academic Press, San Diego, 1989, pp. 576-77) + 50 $\mu\text{g}/\text{ml}$ arginine. Cells were collected by centrifugation of the culture (8,000xg, 10 min, 20°C) and resuspended in a small volume of TAP to give a final density of 2.8×10^8 cells/ml.

In a small sterile test tube containing 0.3 gram of sterile glass beads (0.45 - 0.52 mm), 0.3 ml of this cell suspension, 0.5-1.0 μg of plasmid or 1-2 μg of library DNA, and 0.1 ml of 20% (w/v) polyethylene-glycol (PEG) were added, mixed gently, then vortexed at high speed for 15 seconds using a vortex mixer. The tube was allowed to sit for 2 min and then vortexed for another 15 sec in the same manner.

The cell suspension was then plated, 0.2 ml per plate, onto 2 plates of solid medium (composition: a) when using the arginine auxotroph as a transformation marker: TAP medium + 1.5% (w/v) agar; or b) when using resistance to porphyric herbicides as a transformation marker: TAP medium + 0.1 μM compound A + 50 $\mu\text{g}/\mu\text{l}$ arginine + 1.5% (w/v) agar) and allowed to form colonies under illumination.

C. Transformation by the particle gun method

The particle gun transformation protocol, when employed, followed that described by Boynton, J. E. & Gillham, N. W. (Methods in Enzymol.: Recombinant DNA, Part H, Wu, R. ed., Academic Press, San Diego, CA, 1993, vol. 217, p. 510). The actual protocol employed is described below.

First, the unicellular green alga *Chlamydomonas reinhardtii* strain CC-48 (arginine auxotroph *arg-2*) was cultured mixotrophically for 2 days in TAP liquid medium

composed of 7 mM NH_4Cl , 0.4 mM $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.34 mM $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 25 mM potassium phosphate - 0.5 mM Tris (pH 7.0), 1 ml/L Hutner trace elements, 1 ml/L glacial acetic acid (described in Harris, E. H., The Chlamydomonas Sourcebook, Academic Press, San Diego, 1989) + 50 $\mu\text{g}/\mu\text{l}$ arginine to a cell density of $1.5\text{-}3 \times 10^6$ cells/ml. Cells were collected by centrifugation of the culture (8,000xg, 10 min, 20°C) and resuspended in a small volume of HS medium (composed of 500 mg/L NH_4Cl , 20 mg/L $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 10 mg/L $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 1,440 mg/L K_2HPO_4 , 720 mg/L K_2HPO_4 , 1.2 g/L anhydrous sodium acetate, 1 ml/L Hutner trace elements (described in Harris, E. H., The Chlamydomonas Sourcebook, Academic Press, San Diego, 1989) to a cell density of 1.14×10^8 cells/ml. One ml of this cell suspension was added to small test tubes already containing 1 ml of HS medium + 0.2% agar (Difco Bacto Agar) prewarmed to 42°C. After gentle mixing, 0.7 ml of the suspension was plated onto each of two plates of HSHA agar (composed of 500 mg/L NH_4Cl , 20 mg/L $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 10 mg/L $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 1,440 mg/L K_2HPO_4 , 720 mg/L K_2HPO_4 , 2.4 g/L anhydrous sodium acetate, and 1 ml/L Hutner trace elements (described in Harris, E. H., The Chlamydomonas Sourcebook, Academic Press, San Diego, 1989) also containing 50 $\mu\text{g}/\mu\text{l}$ ampicillin, and the cells were affixed to the surface of the plates by drying them in the dark.

Next, 60 mg of gold particles (1 μm diameter, DuPont Biotechnology Systems, 7556, Wilmington, DE) and 1 ml of ethanol were added to a microtube and vortexed at the highest speed for 2 min using a vortex mixer. The gold particles were subsequently recovered by centrifugation (10,000 rpm, 1 min, room temperature), and this washing procedure was repeated 3 times. The recovered gold particles were subsequently resuspended in 1 ml of sterile distilled water. The particles were again recovered by the same centrifugation procedure, and this washing procedure was repeated another 2 times.

Finally, the gold particles were resuspended in 1 ml of sterile distilled water. Fifty μ l of this particle suspension were added to a microtube, to which 5 μ l of DNA (1 μ g/ μ l), 50 μ l of 2.5M CaCl_2 , and 20 μ l of 0.1M spermidine were added while agitating with a vortex mixer. Mixing was continued for 3 min after which the precipitate was recovered by centrifugation (10,000 rpm, 10 min, room temperature). The precipitated gold particles were resuspended in 250 μ l ethanol, recovered again by the same centrifugation procedure, and finally resuspended in 60 μ l ethanol. *Chlamydomonas* cells prepared as described above were bombarded with the DNA-coated gold particles thus obtained using a particle gun. Immediately afterwards, the cells were recovered from the surface of the agar plates into 1.5 ml of HS liquid medium by scraping gently with a glass rod. Half of this suspension was spread onto each of two selective agar media plates (composition: a) when using the arginine auxotroph as a transformation marker: TAP medium + 1.5% (w/v) agar; b) when using resistance to porphyrin-accumulating type herbicides as a transformation marker: TAP medium + 0.3 μ M compound A + 50 μ g/ μ l arginine + 1.5% (w/v) agar) and allowed to form colonies under illumination ($75\text{--}90\text{ }\mu\text{Molm}^{-2}\text{sec}^{-1}$).

The experimental methods described above were used to screen the genomic DNA library. Details of the screening procedures are presented below as separate primary, secondary, and tertiary screening steps.

1. Primary screening

The recipient unicellular green alga *Chlamydomonas reinhardtii* strain CC-425 (arginine auxotroph *arg-2*, cell wall deficient *cw-15*) was transformed with pARG 7.8 (plasmid DNA) together with the library DNA (a mixture of DNAs extracted from 48 clones) using the glass bead method (see above for details). Half of the cells in each transformation experiment (3.0×10^7 cells) were

used to determine the transformation frequency as indicated by the arginine prototroph phenotype. The remaining half (3.0×10^7 cells) were examined for acquired resistance to porphyric herbicide. This
5 experiment was repeated 198 times, and in total, 9,504 individual clones of the library were screened. In total, 7,046 arginine prototrophs were obtained from 5.8×10^9 cells screened. Assuming all these arginine prototroph colonies are true transformants, the
10 transformation frequency averaged 1.2×10^{-6} . Additionally, one clone was obtained that exhibited resistance to porphyric herbicide (i.e., that grew in the presence of compound A) from 5.8×10^9 cells screened. This colony was also able to grow normally on
15 media lacking arginine, and exhibited a loss of motility when cultured in liquid media.

The DNA pool of 48 clones containing the cosmid which had given rise to the colony exhibiting resistance to porphyric herbicide (cosmid clone 2953 - 3000) is
20 referred to as Cos2953 - Cos3000.

2. Secondary screening

The recipient unicellular green alga *Chlamydomonas reinhardtii* strain CC-48 (arginine auxotroph *arg-2*) was transformed with the DNAs shown in Table 1 by the
25 particle gun method (see above for details). The results are also shown in Table 1.

Transformation with the DNA pool containing the 24 clones Cos2953 - Cos2976 gave rise to colonies resistant to compound A, thus indicating that the porphyrin-
30 accumulating type herbicide resistance gene must be contained within this pool.

Table 1

Sample DNA	No. of colonies exhibiting arginine prototrophy	No. colonies exhibiting resistance to herbicide (per 10 ⁸ cells)
no DNA	0	0
pARG 7.8	165	0
pARG 7.8 + Cos2953-Cos3000	46	4
10 pARG 7.8 + Cos2953-Cos2976	67	20
pARG 7.8 + Cos2977-Cos3000	40	0
pARG 7.8 + Cos5833-Cos5856	29	0
pARG 7.8 + Cos1033-Cos1056	34	0

3. Tertiary screening

15 The recipient unicellular green alga *Chlamydomonas reinhardtii* strain CC-48 (arginine auxotroph *arg-2*) was transformed with hybrid cosmid DNA prepared as described from the respective clones which make up the DNA pool Cos2953 - Cos2976 by the particle gun method (see above

20 for details). These results are shown in Table 2. Transformation only with the hybrid cosmid contained within clone Cos2955 gave rise to colonies resistant to compound A.

Table 2

	Cosmid clone No.	No. of colonies exhibiting resistance to herbicide (per 1.6×10^8 cells)	Cosmid clone No.	No. of colonies exhibiting resistance to herbicide (per 1.6×10^8 cells)
5	No DNA	0	2965	0
	2953	0	2966	0
	2954	0	2967	0
10	2955	28	2968	0
	2956	0	2969	0
	2957	0	2970	0
	2958	0	2971	0
	2959	0	2972	0
15	2960	0	2973	0
	2961	0	2974	0
	2962	0	2975	0
	2963	0	2976	0
	2964	0		

20 In order to confirm this result, purified hybrid
cosmid DNA from Cos2955 was prepared using either a
minicolumn plasmid purification method (Quiagen Inc.) or
the cesium chloride density gradient centrifugation
method, and the transformation experiments were repeated
25 using the same protocol described above. These results
are shown in Table 3. Transformation with Cos2955 DNA
reproducibly gave rise to numerous colonies exhibiting
resistance to compound A, indicating that a porphyrin
herbicide resistance gene must be contained within this
30 hybrid cosmid DNA.

Table 3

Cosmid clone No.	No. of colonies exhibiting resistance to herbicide (per 2.4×10^8 cells)
No DNA	0
2955 (column-purified)	91
SuperCos-1 vector	0
2955 (CsCl purified)	264

10

Example 3Construction of Restriction Map and Subcloning

Hybrid cosmid DNA from clone Cos2955 was purified by the CsCl density gradient centrifugation method. The purified hybrid cosmid DNA (referred to below as Cos2955 DNA), or its subclone isolated as described below, was digested with the restriction enzymes EcoRI, SalI, BamHI, ClaI, KpnI, XhoI, PstI, and HindIII either alone or in combination, and the sizes of the fragments thus generated were estimated by 0.8% agarose gel electrophoresis (25V, 15 hr). From an analysis of the sizes of each fragment in single and double digests, a restriction map was constructed. This result is shown in Figure 1(e). PstI sites were determined in the 3.4 kb DNA fragment shown in Figure 1(a). Cos2955 DNA contains sites for the following restriction enzymes (in order and with the distances (kB) between sites given in parentheses): ClaI (4.4) BamHI (3.1) BamHI (6.6) BamHI (8.2) ClaI (3.1) EcoRI (1.4) XhoI (0.6) PstI (0.5) PstI (0.1) PstI (0.4) PstI (0.1) PstI (0.5) BamHI (0.1) SalI (0.2) SalI (0.9) XhoI (5.1) HindIII (2.8) BamHI (0.2) SalI (0.8) and HindIII. The total molecular size (nucleic acid length) of the nuclear DNA fragment in Cos2955 conferring resistance to porphyrin-accumulating

type herbicides is approximately 40.4 kb.

Cos2955 and the commercially-available plasmid pBluescript-II KS+ (pBS, Stratagene, Inc.) were cut with individual restriction enzymes or an appropriate combination of two restriction enzymes, extracted with phenol/chloroform, and the DNA fragments were recovered by ethanol precipitation. The pBS vector was dephosphorylated by treatment with CIAP if necessary and the pBS vector and the digested Cosmid 2955 DNA fragments were ligated using T4 DNA ligase. The hybrid plasmids thus obtained were introduced into the *E. coli* strain XL1-Blue by electroporation (12.5 kV/cm, 4.5 ms) and spread onto LB agar plates (composed of 10g/L NaCl, 10 g/L Tryptone, 5 g/L yeast extract, 1.5% (w/v) agar and also containing 1 mM IPTG and 50 µg/ml ampicillin) upon which 2% (w/v) X-gal had been spread. From these, white colonies were isolated, i.e., those clones that had taken up the pBS vector and were thus ampicillin-resistant, which had a DNA fragment derived from Cos2955 DNA inserted into the cloning site of the pBS vector and were thus white in color. The isolated colonies were cultured in the presence of ampicillin, and plasmid DNA was subsequently isolated from these colonies by the alkaline lysis method (J. Sambrook, E. F. Frisch, T. Maniatis, Molecular Cloning, 2nd edition, Cold Spring Harbor, Publications, (1989), Vol. I, pp. 1.38 - 1.39). The isolated plasmids were re-digested with the restriction enzyme(s) used for cloning to release the inserts, and the sizes of the inserted fragments obtained were again estimated by 0.8% (w/v) agarose gel (75V, 5 hr) electrophoresis. When an insert of the desired size was obtained, it was subjected to further restriction analysis in order to confirm that the correct DNA fragment had been cloned. The DNA fragments thus cloned are shown in Figures 1(a)-1(d). In order to identify the clone containing the porphyrinic herbicide resistance mutation *rs-3*, the recipient *Chlamydomonas reinhardtii* strain

CC-48 (arginine auxotroph *arg-2*) was transformed with DNA from the pBS subclones of Cos2955 by the particle gun method (see above for details). The pBS subclones of Cos2955 that were able to confer resistance to compound A contained the Eco13.8, Hind10.0, and Xho3.4 fragments. These results confirmed that these DNA fragments contain the porphyrin herbicide resistance mutation *rs-3*.

Herbicide-resistant transformants were obtained from the 13.8 and 10.0 kb fragments from RS-3 at about 80-fold higher frequency than from the 3.4 kb fragment (Table 4). This is consistent with the 13.8 and 10.0 kb fragments containing the entire coding sequence plus upstream and downstream regulatory elements, and integrating non-homologously and randomly in the nuclear genome of the herbicide-sensitive recipient strain. In contrast, the low transformation frequency observed with the 3.4 kb fragment is consistent with this fragment containing only a portion of the RS-3 gene which must integrate into the herbicide-sensitive RS-3 gene of the recipient by homologous recombination to be expressed. Nuclear transformants of *Chlamydomonas reinhardtii* arise much more frequently by random non-homologous recombination than by homologous recombination, as has been demonstrated for the nuclear *nit-1* gene by Sodeinde, O.A. and Kindle, K.L. (Proc. Natl. Acad. Sci. USA vol. 90, p. 9199 (1993)).

Table 4

pBS insert	No. of colonies exhibiting resistance to herbicide (per 2.4×10^8 cells)
No DNA	0
5 Eco13.8 or Hind10.0	894
Xho3.4	10

Eco13.8 is a DNA fragment of approximately 13.8 kb that is able to confer resistance to porphyric herbicides, and which contains sites for the following restriction enzymes (in order and with the distance (Kb) between sites given in parentheses; this same notation will be used throughout): EcoRI (1.4) XhoI (0.6) PstI (0.5) PstI (0.1) PstI (0.4) PstI (0.1) PstI (0.5) BamHI (0.1) SalI (0.2) SalI (0.9) XhoI (5.1) HindIII (2.8) BamHI (0.2) SalI (0.8) and HindIII (0.1>) KpnI. The restriction site map for this fragment is shown in Figure 1(d). Hind10.0 is a DNA fragment of approximately 10.0 kb that is able to confer resistance to porphyric herbicides, and which contains sites for the following restriction enzymes: EcoRI (1.4) XhoI (0.6) PstI (0.5) PstI (0.1) PstI (0.4) PstI (0.1) PstI (0.5) BamHI (0.1) SalI (0.2) SalI (0.9) XhoI (5.1) HindIII (0.1>) KpnI. Its restriction site map is shown in Figure 1(c). Furthermore, the Hind10.0 fragment is a derivative of the Eco13.8 fragment listed above from which has been deleted a DNA fragment of approximately 3.8 kb containing sites for the restriction enzymes HindIII (2.8) BamHI (0.2) SalI (0.8) HindIII. Hind9.9 (Figure 1(b)) is derived from Hind10.0 by deletion of an

approximately 0.1 kB fragment (see Figures 1(b) and 1(c)). Xho3.4 is a DNA fragment of approximately 3.4 kB that is able to confer resistance to porphyric herbicides, and which contains sites for the restriction enzymes XhoI (0.6) PstI (0.5) PstI (0.1) PstI (0.4) PstI (0.1) PstI (0.5) BamHI (0.1) SalI (0.2) SalI (0.9) XhoI. Its restriction site map is shown in figure 1(a). Xho3.4 is a derivative of Eco13.8, described above, from which has been deleted a DNA fragment of approximately 9.1 kB containing sites for the restriction enzymes XhoI (5.1) HindIII (2.8) BamHI (0.2) SalI (0.8) HindIII (0.1) KpnI and a DNA fragment of approximately 1.4kB containing sites for the restriction enzymes EcoRI (1.4) XhoI.

E. coli strains containing pBS plasmids with the fragments described above inserted, i.e., Eco13.8, Hind10.0, and Xho3.4 (Figures 2-4) have been deposited with the *Chlamydomonas* Genetics Center, c/o Dr. Elizabeth H. Harris, DCMB Group, LSRC Building, Research Drive, Box 91000, Duke University, Durham, North Carolina, 27708-1000. These deposits are designated P-563, P-564, and P-566, respectively. *E. coli* containing Cos2955 has also been deposited with the *Chlamydomonas* Genetics Center under the designation P-561. In addition, *Escherichia coli* XL1-BLUE/Eco13.8 has been deposited with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD, 20852, USA, on July 19, 1995 under the terms of the Budapest Treaty, and has been given the deposit designation ATCC 69870.

Example 4

Determination of the Nucleotide Sequence of the Xho3.4 DNA Fragment

The nucleotide sequence of the Xho3.4 DNA fragment obtained as described in Example 3 was determined by the Sanger enzymatic sequencing method (Sequenase Version

2.0 kit, USB Inc.) using α -³⁵S-dATP label. This protocol is described below.

Approximately 5 μ g of the Xho3.4 DNA fragment obtained as described in Example 4, a plasmid containing this fragment, or a plasmid containing the 1.7 kb PstI-XhoI fragment of the Xho3.4 fragment was incubated in 0.2M NaOH, 0.2M EDTA at 37°C for 30 min. The DNA was recovered by ethanol precipitation and washed with 70% (v/v) ethanol. To this precipitate were added 7 μ l of deionized distilled water, 2 μ l of Sequenase annealing buffer (5-fold concentration), 1 μ l of primer to give a final composition of 40 mM Tris-HCl (pH 7.5), 20 mM MgCl₂, 50 mM NaCl, and 5 μ g/L primer. This solution was heated to 65°C for 2 min, and allowed to cool to room temperature below 30°C over 30 min. The solution was then placed on ice. To this mixture were added 1 μ l of 0.1M DTT (dithiothreitol), 2 μ l of labeling mixture (containing 1.5 μ M dGTP, 1.5 μ M dCTP, 1.5 μ M dTTP), 0.5 ml of [α -³⁵S]dATP (10 μ Ci/ml, 1000 Ci/mmol), and 3 units of Sequenase version 2.0 T7 DNA polymerase. The solution was mixed well and allowed to react at room temperature for approximately 5 min. At the same time, four microtubes were individually labeled G, A, T, and C, and to these were added 2.5 μ l of the respective ddNTP reaction solutions (composed of 80 μ M dGTP, 80 μ M dATP, 80 μ M dTTP, 80 μ M dCTP, 50 mM NaCl, and 8 μ M of either dideoxy-dGTP, dideoxy-dATP, dideoxy-dTTP, or dideoxy-dCTP). These tubes were pre-incubated at 37°C for 1 hr. Three point five μ l of the completed labeling reaction mixture were then added to each of these four tubes, quickly mixed, and allowed to react for approximately 5 min at 37°C. The reaction was halted by the addition of 4 μ l of reaction stop solution (containing 95% formamide, 20 mM EDTA, 0.05% bromophenol blue, and 0.05% xylene cyanol FF) and immediately mixed. The samples were analyzed by 8M urea-6% (w/v) polyacrylamide gel electrophoresis (approximately 65 watts,

approximately 4 hr), the gel dried and exposed to Kodak XAR-5 film, and the sequence determined from the resultant autoradiogram. The resulting nucleotide sequence is shown in SEQ ID NO:1.

5

Example 5

Biologically Functional Equivalent DNA Fragments
Conferring Resistance to Porphyrin-Accumulating
Type Herbicides

Each of the nucleic sequences disclosed herein, or
10 their biologically functional equivalents, can be used
in accordance with the present invention. The phrase
"biologically functional equivalents," as used herein,
denotes nucleic acid sequences exhibiting the same or
similar biological activity as the particular nucleic
15 acid sequences described herein, i.e., when introduced
into plant or algal cells in a functionally operable
manner so that they are expressed, they confer
resistance to porphyrin-accumulating type herbicides
thereon.

20 For example, the nucleic acid sequences described
herein can be altered by base substitutions, additions,
or deletions to produce biologically functionally
equivalent nucleic acids that encode proteins conferring
resistance to porphyrin herbicides *in vitro* and *in vivo*.
25 In addition, due to the degeneracy of the genetic code,
other DNA sequences that encode substantially the same
amino acid sequences as described herein and confer
resistance to porphyrin herbicides *in vitro* and *in vivo*
can be used in the practice of the present invention.
30 These include, but are not limited to, nucleotide
sequences comprising all or portions of the genomic DNAs
described herein or the corresponding mRNAs or cDNAs
that are altered by the substitution of different codons
that encode a physiologically functionally equivalent
35 amino acid residue within the protein sequence, thus
producing a silent change. Similarly, the proteins

conferring porphyric herbicide resistance, or derivatives thereof, encoded by the present invention include, but are not limited to, those containing all of the amino acid sequences encoded by the DNA sequences substantially as described herein, including altered sequences in which functionally equivalent amino acid residues are substituted for residues within the sequence, resulting in a silent change. For example, one or more amino acid residues within the sequence can be substituted with another amino acid of similar polarity which acts as a functional equivalent, resulting in a silent alteration. Substitutes for an amino acid within the sequence may be selected from other members of the class to which the amino acid belongs. For example, fungible nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine. Fungible polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine. Fungible positively charged (basic) amino acids include arginine, lysine, and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid.

The variants of the genomic DNAs, the corresponding mRNAs or cDNAs, and proteins contemplated herein should possess more than 75% homology, preferably more than 85% homology, and most preferably more than 95% homology, to the naturally occurring genomic DNAs, the corresponding mRNAs or cDNAs, and proteins discussed herein. To determine this homology, two proteins (or nucleic acids) are aligned so as to obtain a maximum number of matched residues using gaps and inserts. Homology (of two proteins) is determined as the result of the number of matched amino acids divided by the number of total amino acids plus gaps and inserts, multiplied by 100.

Biologically functional equivalents to the nucleic

acid fragments disclosed herein can be created by mutagenesis techniques such as those described, for example, in Osuna, J., Flores, H., and Soberon, X. (Critical Reviews in Microbiology 20:107-116 (1994)), and selected for by transformation and screening of *Chlamydomonas* as described in Example 2.

Example 6

Isolation of DNA Fragments That Confer Resistance to Porphyrin-Accumulating Type Herbicides From Organisms Other Than *Chlamydomonas reinhardtii*

Degenerate oligonucleotide primers based on the deduced amino acid sequence of a cDNA corresponding to the RS-3 gene from *Chlamydomonas reinhardtii* can be synthesized and used to isolate the equivalent gene or cDNA from eubacteria, cyanobacteria, algae, and higher plants. PCR technology employing reverse transcriptase (Kawasaki, E.S. PCR Protocols, Ch. 3, p. 21, Innis et al., eds., Academic Press, San Diego, (1990)) and these degenerate oligonucleotides can be used to amplify the equivalent cDNAs from algae and higher plants, and these can be easily cloned in appropriate transformation/expression vectors for dicots such as pBIN described by Bevan, M. (Nucl. Acids Res. Vol. 12, p. 8711 (1984)) or pZ597 as described by Svab, Z. et al. (Plant Mol. Biol. Vol. 14, p. 197 (1990)), or for monocots using pDB1 described by Becker, D. et al. (Plant J., Vol. 5, p. 299 (1994)) or pBARGUS described by Vasil, V. et al. (Bio/Technology, Vol. 10, p. 667 (1992)). Transformed plants expressing the cDNAs in these vectors can be isolated using established procedures as described below. Alternatively, the degenerate oligonucleotides can be used as probes to screen cDNA libraries from crop plants in lambda phage vectors such as λ ZapII (Stratagene). The corresponding cDNA can be transferred to an appropriate transformation/expression vector for introduction into monocot or dicot crop plants as

described below. In the case of eubacteria and cyanobacteria, the degenerate oligonucleotides can be used to screen genomic libraries directly, and the appropriate coding sequences can be transferred into one of these transformation/expression vectors for crop plants as described below.

Example 7

Transformation of Crop Plants and Algae to Porphyrin-Accumulating Type Herbicide Resistance

The research described herein has identified DNA fragments that are able to confer resistance to porphyrinic herbicides onto plants and algae. Crop plants can be made resistant to porphyrinic herbicides by the introduction therein of these DNA fragments or their biologically functional equivalents. This invention will permit the use of porphyrinic herbicides during crop cultivation, and thus facilitate weed control and cultivation management on such crops.

Since the *Chlamydomonas rs-3* resistance mutation is dominant (Sato, R. et al., ACS Symposium Series, Vol. 559, Porphyrinic Pesticides, S.O. Duke and C.A. Rebeiz, eds., Chapter 7, p. 91 (1994)), a full length cDNA corresponding to the subject DNA fragments under the control of the appropriate upstream and downstream regulatory sequences can be introduced into crop plants that lack resistance to porphyrin-accumulating type herbicides, so the RS-3 resistance gene can be utilized to generate crop plants resistant to porphyrinic herbicides.

In the case of dicot crop species, the chimeric RS-3 resistance construct can be inserted into a binary *Agrobacterium* vector such as pBIN described by Bevan, M. (Nucl. Acids Res. Vol. 12, p. 8711 (1984)) or pZ597 as described by Svab, Z. et al. (Plant Mol. Biol. Vol. 14, p. 197 (1990)), transferred into *Agrobacterium* by triparental mating according to Hoekema, A. et al.

(Nature Vol. 303, p. 197 (1983)), and transformed into leaf discs of an agriculturally desirable cultivar by co-cultivation on medium containing carbenicillin and kanamycin. Shoots can be regenerated from kanamycin-resistant calli, and induced to form roots using standard plant tissue culture protocols (Bevan, M. Nucl. Acids Res. Vol. 12, p. 8711 (1984)). Kanamycin-resistant plants can be assayed for levels of resistance to porphyric herbicides, and the expected dominant 3:1 segregation of resistance verified in crosses to sensitive cultivars. Depending on the breeding system for a particular crop species, the herbicide-resistant transformants can be propagated by self pollination or backcrossed to the sensitive cultivar to establish a pure breeding, herbicide-resistant line. Examples of dicotyledonous crop plants to which this method can be applied include alfalfa, beans, cabbage, carrots, clover, cotton, various cucurbits, flax, peas and other agronomically important legumes, peanuts, peppers, potatoes, soybeans, sugar beets, sunflower, tobacco, and tomatoes.

In the case of monocot crop species, the full length RS-3 cDNA can be inserted in a monocot expression vector such as pDB1 described by Becker, D. et al. (Plant J., Vol. 5, p. 299 (1994)) or pBARGUS described by Vasil, V. et al. (Bio/Technology, Vol. 10, p. 667 (1992)) fused to the *Adh1* promoter and intron or the *Act1* promoter and to the *nos* terminator replacing the GUS gene in the constructs. Depending on the species, the chimeric plasmid can be introduced into embryogenic calli, immature embryos, scutellar tissue, immature inflorescences, microspores, or protoplasts by biolistic transformation, and calli, shoots, and plants regenerated under selective conditions in the presence of glufosinate to which the *bar* gene product encoded by both plasmids confers resistance. Glufosinate-resistant transgenic plants can then be assayed for levels of

resistance to porphyric herbicides, and the expected dominant 3:1 segregation verified in crosses to sensitive cultivars. Depending on the breeding system for a particular crop species, the herbicide-resistant transformants can be propagated by self pollination or backcrossed to the sensitive cultivar to establish a pure breeding, herbicide-resistant line. Examples of monocotyledonous crop plants to which this method can be applied include barley, corn, forage crops, oats, onions, rice, rye, sorghum, sugar cane, and wheat.

If such crop plants that have thereby acquired resistance to porphyric herbicides are cultivated, the utilization of these herbicides on these crop plants becomes feasible. This should allow for simpler and more effective weed management, and increase the value of these herbicides in agricultural use. Furthermore, by using the subject DNA fragments as probes, it should be possible to identify other DNA fragments in crop plants that exhibit a high degree of sequence homology to the subject DNA fragments. This should make it possible to assess qualitatively and/or quantitatively whether a given crop plant will be resistant to porphyric herbicides prior to their actual treatment with such herbicides. In addition, this gene could be used as a resistance-type genetic marker in plant genetic engineering research and plant molecular biology/biotechnology, and should thus have significant industrial application.

The invention being thus described, it will be obvious that the same may be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the invention, and all such modifications as would be obvious to one skilled in the art are intended to be included within the scope of the following claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Sato, Ryo
Boynton, John E.
Gillham, Nicholas W.
Harris, Elizabeth H.
- (ii) TITLE OF INVENTION: Porphyrin Accumulating-Type Herbicide Resistance Gene
- (iii) NUMBER OF SEQUENCES: 1
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Birch, Stewart, Kolasch & Birch, LLP
 - (B) STREET: P.O. Box 747
 - (C) CITY: Falls Church
 - (D) STATE: Virginia
 - (E) COUNTRY: USA
 - (F) ZIP: 22040-3487
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: PT New
 - (B) FILING DATE: 19-JUL-1996
 - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: PT PCT/US95/09098
 - (B) FILING DATE: 20-JUL-1995
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Murphy Jr., Gerald M.
 - (B) REGISTRATION NUMBER: 28,977
 - (C) REFERENCE/DOCKET NUMBER: 2185-154F(PC)
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 703-205-8000
 - (B) TELEFAX: 703-205-8050

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 3381 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: not relevant
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: YES
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Chlamydomonas reinhardtii
 - (B) STRAIN: RS-3

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CTCGAGTTGG CATTGTTAGT CTTAAGCATC TTCGAAACCA TGGCACTTAT GGGCGAAGCC	60
GCAACATACA CCTGACGATG CTCCCCTGCC CTCCTCTTCC CACCACGGAA GCAACCCATG	120
CAATTGGTCT TTCGTTGTG CACTGACAGT CAATGCCTTG GGTCTGGCCC CCTGCTCCGG	180
GTATTGCGAA TTTATAGCCC CCTTTTGAAG GTACTTCGAG ACCTAGTCTA TGTCACCTCAG	240
TTCTGTGTCT CTCCTGTGCT ATGCTGGGAC CACACTGAGG GACAGTGCCC CCAAACCGCC	300
CCGCGTGTGC GCCAACTCAC TCTCGCCAAA ACTTCATGCA AACAATGCAC AACGGGGCTG	360
TACAACGGAG CGTCATAGTG CAGTTAACCC GCACATAACG GTGCCGACAG AGCTTTACCA	420
ACCACATCGC TCATCCCAGC ACCACCTATG TCCTTGGCAG ACCCCCGAAC CCCTAGCTCC	480
ACATCCCCGA TCCATTCCAA CTGTTACTAC ACATCCCAGC AAGCGCCAAT GGTAGCCCCC	540
CCCTCGCCTC CACTCCGCTT GCCCCAAGTG CTTGCCAATT GCTGCTGCCG GTGCCACTGC	600
GGCATTTACC AGTACAGCC CGATGCTGCT GCTGCTACTG CTGCAGCCGC TTAGGCCTTG	660
ACTGCGGCCT TGGACACGCT CTTGGCCAGG TTGGCTGCGG ACTCGTAGCC GTGCTCCACC	720
ACCTTGCCCA GGGCCACACC TGGTTGCAGG GGTTCGAGGG GGGAGAGGGG CCGAGGGGGT	780
TGGGGATGAG GCAAAGGCGT GGGCACACGT GTGGGCGGGG GCGGGCTGGG CGGCTGGGCA	840
CCATACAGCG AAGGGGCCAG GGGGGTCATG TGCCCCGAGCC CAACGAAATA GGTCCCAATA	900
GTCAAGGAAT CATCGCCCGC GTCGGGGGTA GTCATACATG TAGCCTCCGC TCCTTGCACT	960
CCCCTGACCT CTTGTTTCATG CTGGGTTTCC CGGTTTCAGT CAATGAGGAT ACAACACCGC	1020
CTTGCTTTGC GCCTTAGCAC CACCTCCTGC TCAACCTCCT CTTGTACCC TCCCCTCCCC	1080
TCCCCTCCCC TCTTCCTGCT GCTGCTGCTG CCCACGCGCT CACCGCTGAC GTAGTTGCCC	1140
CCCAGGTGCA CGCCCTGCAG CCCCGCCGCG TCCAGCGCCT TGCGCGCCTT GTCCAGCTGC	1200
TCCAGGTGGC CCAGGTTGAA CTGCAGGTGG CAGGGCGGGG AGGAGGGGTT ACACATGCAG	1260
CCCAAATAA ACCGAACCTA ATGCAAAGAG TGGTACAGAG AAGTGCAGCA AAGTATACAA	1320
CAGGTGAAGT TGATTCCAGA ACCCAAAAGC AAGTCAACCG AGTGAACCCA AATCCGGGGA	1380
GAGGTGGCTA GGTGGGAGG GGGCAGCCAG GTTATCAGGT CGAAGGTCGC AGGTCGAGGT	1440
TATAGGGCTG TAGAGCTCTT TGAGAAACTA GGATCTAGCC CATCCCTCCG GCTGCTGCGC	1500
CCTCACACCT GCGGGATGGC GCGCGGCCAC ACGCGCACGC CCACCACACG GGGCTTGGGC	1560
GCGTCGGGCT TGATGACCAT GTTGCGCAGG TCCTTGTTCCA CCTGCAGGGA GATGGGGTGG	1620
GGGGAGAGGC GGTGAGAGCA GCGTGCGGC AGGGGGGGGG GTGCGGCAGG TGAAGACGT	1680
GGAACGGGCG GCGGGGTTG GGGCTGCAGG ACGAAAATGT GTGTGTGTGT GTGTGTGTGT	1740
GTGTGTGTGT GTGTGTGTGT GTGTGTGTGT GTGTGTGTGT GTGTGTGTGT GTGTGTGTGT	1800
GTGTGTGTGT GTGTGTGTGT GTGTGTGTGT GTGTGTGTGT GTGTGTGTGT GTGTGTGTGT	1860
GAAGCAACAC GGGTCCAAGA CGCCACACAA CAAGGGGTGT CAGCTTTGTA CAGCTGGAAA	1920
ATGGGCTGAG CCCCATGTAG CGCAAGGGGA TGTGACGATG GGCACCGGCG ACATGAACAG	1980

CACCCTTGCA AGCCCATCCA GCGGGACTGG CCACTAGGCC AACGCTGTGC CGAGGCACCG	2040
ACATGCCCTG TCCCCGCCAC GCCACGCCAC CTTACCCACG CCACTTCACG GTCCATGATA	2100
TCCCAAATGC ACCTCACCCA CATCCCCTC AAGCACCACA AAGGGCTAGC CGGGCTCGGG	2160
CGCGGGATCC CGGGCCGCGA CACATGTAAG GCTCGGGGAC GCATGGTTTG GCTGTTGCAA	2220
ACAATTTTGA CATTCTTGCC CCAAGACGCT TGTCGTCGAC ATGTTTTGAT ACATGGATGT	2280
AAGATATTTA GGGGCCGAGA GCTATACTCG CGAACTGAAG AAAGTCAAGA TGTCCATGGA	2340
CTCACGAGGT CGCTTCTCGC TCCGGCCGAG TTTTCCCTGC CGTCTATTTT TCTACAATAG	2400
GTAATTGCAA TTAATAATAC GTGGCACTCG CGGTTTGTGC CAGTACGGTA CTTTTTGTGC	2460
ACACAGGCAC ACACGCACGC ACGGACAGGG GCAATCCTGG GGGGCTTCGC CCCCCCTGGA	2520
TCGCTTTGCT CAGGGGGCTC AGCCCAAAAA TCCACTGCCC CCCCCCACC CCCCCACACA	2580
CACACCTGCT CCACCAGCTG CTCGGTGGTC TGGTTGACGA TGCCGCGGTT GGTGGTGCCG	2640
CCGATGTAGT TGAGCAGCAG CATGTGGCCC TCGGGCGCGC GGCCGGGGAA CAGGCTGGAG	2700
CTGTAGATGG TGCCAGAGT GGTGATGCC TGTGTGTGGA CAAGTGTGTG TGTTAGAAGA	2760
CACCAAAATG AAGCGAAGAG TGTGTTAAGG AGCACCAGAC AAAGTAAGCG CAAAGAGGGT	2820
GCGTGTGCGC CCGTGTGTGT TGTGAAGGG GGGAAATGGAT GAGGGGAGCG CGAGGATACA	2880
ACCGCGGGAT ACGACCCAGC GCCCCAATCC CCCCCACCC CACCCCCAAC CCCCACCACC	2940
CTCCCCCTCG CAACACACCG CCCGCAACAC GCGCGCACTT GCCCACCTGC GTGCGCGGGT	3000
GCAGCTGACC GAAGCCCGGC ACGGACCCGT CCGAGGCCTT GCGCTCCTCC CGCACGGCGC	3060
TCAGCGGGTA CGACAGCGTC ACGGCGCCCA TCGGCGGGTA GTCGAAGGAG CCCAGGGCCT	3120
CGGCGGCGGC GGGCTGTGCG GGGGGGAGAG GGAGGGAGGG GCAGGCGCAG GGAGGCGGGG	3180
TTACGTTAAT GATTGCCCAA GAACTGGTA GACGGTAGAC AGTCTAGGTG GGGGAGGAGG	3240
AGCGGATGGA ATCGGGATGG AGCCGAGGAG TGGGAAGGGG AGTAAAGCCG GGGGGGAGCG	3300
GGTAGCAGGA AAGGGGGACG TGGCCGTGCA CACAAAGAAG CCGGAACAGG TGCCAAACGG	3360
ATTTCCTCCA ACGCTCTCGA G	3381

WHAT IS CLAIMED IS:

1. An isolated, purified DNA fragment, having a nucleotide sequence of a portion of the DNA comprising the genome of an alga, or having a nucleotide sequence
5 highly homologous to a nucleotide sequence of a portion of the DNA comprising the genome of an alga, that confers resistance to porphyrin-accumulating type herbicides when expressed in plant or algal cells.
2. A plasmid containing said DNA fragment of claim
10 1.
3. A microorganism containing said plasmid of claim 2.
4. A method for conferring resistance to porphyrin-accumulating type herbicides upon plant or
15 algal cells, comprising introducing said DNA fragment of claim 1 into said plant or algal cells, wherein said DNA fragment is expressed.
5. A plant or alga into which said DNA fragment of claim 1 has been introduced, wherein said DNA fragment
20 is expressed.
6. An isolated, purified DNA fragment having the

following characteristics:

- a) comprising a nucleotide sequence of a DNA fragment obtained from a strain of the unicellular green alga *Chlamydomonas reinhardtii* that exhibits resistance to porphyrin-accumulating type herbicides;
 - b) containing restriction sites for XhoI, PstI, PstI, PstI, PstI, BamHI, SalI, SalI, and XhoI, and having a restriction site map as shown in Figure 1(a);
 - c) having a molecular size of approximately 3.4 kb;
- and
- d) which confers resistance to porphyrin-accumulating type herbicides in plant or algal cells when expressed therein;
- or a biologically functional equivalent of said DNA fragment.

7. An isolated, purified DNA fragment having the following characteristics:

- a) comprising a nucleotide sequence of a DNA fragment obtained from a strain of the unicellular green alga *Chlamydomonas reinhardtii* that exhibits resistance to porphyrin-accumulating type herbicides;
 - b) containing restriction sites for EcoRI, XhoI, PstI, PstI, PstI, PstI, PstI, BamHI, SalI, SalI, XhoI and HindIII, and having a restriction site map as shown in Figure 1(b);
 - c) having a molecular size of approximately 9.9 kB;
- and
- d) which confers resistance to porphyrin-accumulating type herbicides in plant or algal cells when expressed therein;
- or a biologically functional equivalent of said DNA fragment.

8. An isolated, purified DNA fragment having the following characteristics:

- a) comprising a nucleotide sequence of a DNA

fragment obtained from a strain of the unicellular green alga *Chlamydomonas reinhardtii* that exhibits resistance to porphyrin-accumulating type herbicides;

5 b) containing restriction sites for EcoRI, XhoI, PstI, PstI, PstI, PstI, PstI, BamHI, SalI, SalI, XhoI, HindIII, and KpnI, and having a restriction site map as shown in Figure 1(c);

 c) having a molecular size of approximately 10.0 kb; and

10 d) which confers resistance to porphyrin-accumulating type herbicides in plant or algal cells when expressed therein; or a biologically functional equivalent of said DNA fragment.

15 9. An isolated, purified DNA fragment having the following characteristics:

 a) comprising a nucleotide sequence of a DNA fragment obtained from a strain of the unicellular green alga *Chlamydomonas reinhardtii* that exhibits resistance to porphyrin-accumulating type herbicides;

20 b) containing restriction sites for EcoRI, XhoI, PstI, PstI, PstI, PstI, PstI, BamHI, SalI, SalI, XhoI, HindIII, BamHI, SalI, HindIII, and KpnI, and having a restriction site map as shown in Figure 1(d);

25 c) having a molecular size of approximately 13.8 kb; and

 d) which confers resistance to porphyrin-accumulating type herbicides in plant or algal cells when expressed therein; or a biologically functional equivalent of said DNA fragment.

30 10. A plasmid containing said DNA fragment or biologically functional equivalent of said DNA fragment of any one of claims 6, 7, 8, or 9.

11. A microorganism containing said plasmid of claim 10.

12. A method of conferring resistance to porphyrin-accumulating type herbicides upon plant or algal cells, comprising introducing said DNA fragment or biologically functional equivalent of said DNA fragment of any one of claims 6, 7, 8, or 9 into said plant or algal cells in a functionally operable manner so that said DNA fragment or biologically functional equivalent of said DNA fragment is expressed in said plant or algal cells.

13. A plant or alga into which has been introduced in a functionally operable manner said DNA fragment or biologically functional equivalent of said DNA fragment of any one of claims 6, 7, 8, or 9.

14. An isolated, purified DNA fragment comprising the nucleotide sequence shown in SEQ ID NO:1.

15. A plasmid containing said DNA fragment of claim 14.

16. A microorganism containing said plasmid of claim 15.

17. A method of conferring resistance to porphyrin-accumulating type herbicides upon plant or algal cells, comprising introducing cDNA corresponding to mRNA encoded by said DNA fragment of claim 14 into said plant or algal cells, wherein said cDNA is expressed.

18. A plant or alga into which cDNA corresponding to mRNA encoded by said DNA fragment of claim 14 has been introduced, wherein said cDNA is expressed.




FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4

**CORRECTED
VERSION**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/11, 15/70, 1/12, 1/21, 15/82, A01H 5/00	A3	(11) International Publication Number: WO 97/04089 (43) International Publication Date: 6 February 1997 (06.02.97)
(21) International Application Number: PCT/US96/11999 (22) International Filing Date: 19 July 1996 (19.07.96) (30) Priority Data: PCT/US95/09098 20 July 1995 (20.07.95) WO (34) Countries for which the regional or international application was filed: US et al. (71) Applicants (for all designated States except US): SUMITOMO CHEMICAL COMPANY, LTD. (JP/JP); 5-33, Kitahama 4-chome, Chuo-ku, Osaka 541 (JP). DUKE UNIVERSITY [US/US]; 012 Allen Building, Durham, NC 27708 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): SATO, Ryo (JP/JP); 2-10-5-301 Sonehigashinocho, Toyonaka-shi, Osaka 561 (JP). BOYNTON, John (US/US); 1808 Woodburn Road, Durham, NC 27705 (US). GILLHAM, Nicholas, W. (US/US); 1211 Woodburn Road, Durham, NC 27705 (US). HARRIS, Elizabeth, H. (US/US); 3413 Rolling Hill Road, Durham, NC 27705 (US). (74) Agents: BIRCH, Anthony, L. et al.; Birch, Stewart, Kolasch & Birch, L.L.P., P.O. Box 747, Falls Church, VA 22040-0747 (US).	(81) Designated States: CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 9 May 1997 (09.05.97)	
(54) Title: PORPHYRIN-ACCUMULATING TYPE HERBICIDE RESISTANCE GENE (57) Abstract Provided are DNA fragments and biologically functional equivalents thereof that confer resistance to porphyrin-accumulating type herbicides upon plant and algal cells, plasmids containing these DNA fragments or biologically functional equivalents thereof, microorganisms containing these DNA fragments or biologically functional equivalents thereof, methods for conferring resistance to porphyrin-accumulating type herbicides upon plant or algal cells using these DNA fragments or biologically functional equivalents thereof, and herbicide-resistant plants or algae into which these DNA fragments or biologically functional equivalents thereof have been introduced and in which they are expressed.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/11999

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12N15/11 C12N15/70 C12N1/12 C12N1/21 C12N15/82
A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	AM.CHEM.SOC. SYMPOSIUM SERIES, vol. 559, 1994, pages 91-104, XP000651696 SATO, R. ET AL.: "Characterization of a mutant of Chlamydomonas reinhardtii resistant to protoporphyrinogen oxidase inhibitors" see page 93, paragraph 2 - paragraph 3 see page 95, paragraph 3 - page 96, left-hand column see page 98, last paragraph - page 103, paragraph 2; figure 4 --- -/--	1-18

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

12 March 1997

Date of mailing of the international search report

01. 04. 97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Donath, C

INTERNATIONAL SEARCH REPORT

Int. Application No
PCT/US 96/11999

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>ZEITSCHRIFT FÜR NATURFORSCHUNG, vol. 48, no. 3-4, 1993, pages 339-344, XP000651400 OSHIO, H. ET AL.: "Isolation and characterization of a Chlamydomonas reinhardtii mutant resistant to photobleaching herbicides" see abstract see page 342, left-hand column, last paragraph - page 343, right-hand column, paragraph 2</p>	1-18
A	<p>--- J. PESTICIDE SCIENCE, vol. 15, no. 3, 1990, pages 449-451, XP000651693 KATAOKA, M. ET AL.: "Isolation and partial characterization of mutant Chlamydomonas reinhardtii resistant to herbicide S-23142" see the whole document -----</p>	1-18